

~~SECRET//NOFORN~~

SARS-COV-2 GENOME ANALYSIS


(b)(3); 10 USC 424; (b)(6)

June 25, 2020

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

SARS-CoV-2 Genome



The next two pages are DIF citing (b)(1) and (b)(3)
and are not provided.

~~SECRET//NOFORN~~

Methods to construct a Coronavirus Full-Length Clone

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~


CORONAVIRUS INFECTIOUS CLONE CONSTRUCTION

- 1) Synthesize or PCR amplify 6-10 segments of a Bat CoronavirusBuild a 5' transcription initiation fragment "stitch" the fragments together using an infectious clone technology 3' to the transcription initiation fragmentRestriction-enzyme-based fragment cloning systemOverlapping Fragment systemGuided RNA RecombinationClone in a suitable host (E. coli, yeast, etc.)Sequence verify cloned insert

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

CAPABILITIES NEEDED TO CONSTRUCT A BAT-LIKE CORONAVIRUS INFECTION CLONE



The next page is DIF citing (b)(1) and (b)(3) and is not provided.

~~SECRET//NOFORN~~

Type IIS Restriction Enzymes and
Golden Gate Assembly System

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

Type IIS Restriction Enzymes

- Non palindromic recognition siteCuts at sites outside of recognition siteEach digested location has unique nucleotide overhangsExampleBsal

5' – GGTCTCNNNN – 3'
3' – CCAGAGNNNN – 5'



~~SECRET//NOFORN~~

~~SECRET//NOFORN~~


Eleven Tips For Optimizing Your Golden Gate Assembly Reactions

Looking to make the most DNA fragments in a single reaction? Here are some tips to help you start planning your Golden Gate Assembly.

- 1. Check your enzymes**
Enzymes are the workhorses of your assembly. Make sure you have the right enzymes for the job. For example, if you're using a restriction enzyme, make sure it's compatible with the vector and the other enzymes in the reaction. If you're using a ligase, make sure it's compatible with the vector and the other enzymes in the reaction.
- 2. Optimize your enzymes**
When using multiple enzymes in a reaction, it's important to optimize each one. For example, if you're using a restriction enzyme, make sure it's compatible with the vector and the other enzymes in the reaction. If you're using a ligase, make sure it's compatible with the vector and the other enzymes in the reaction.
- 3. Choose the right plasmid**
The plasmid you choose will affect the efficiency of your assembly. Make sure you choose a plasmid that is compatible with the enzymes you're using. For example, if you're using a restriction enzyme, make sure it's compatible with the vector and the other enzymes in the reaction.
- 4. Choose the right buffer**
The buffer you use will affect the efficiency of your assembly. Make sure you choose a buffer that is compatible with the enzymes you're using. For example, if you're using a restriction enzyme, make sure it's compatible with the vector and the other enzymes in the reaction.
- 5. Increase your complexity**
Complexity is key to increasing the Golden Gate Assembly's efficiency. Make sure you include all the necessary components in your reaction. For example, if you're using a restriction enzyme, make sure it's compatible with the vector and the other enzymes in the reaction.
- 6. Make sure your plasmid prep is RAMP-free**
RAMP-free plasmid prep is essential for a successful Golden Gate Assembly. Make sure you use a RAMP-free plasmid prep kit to ensure that your plasmid is free of contaminants.
- 7. Avoid protein inhibitors**
Protein inhibitors can interfere with the enzymes in your reaction. Make sure you avoid protein inhibitors like SDS and Triton X-100.
- 8. Avoid PCR-induced errors**
PCR-induced errors can lead to inaccurate results. Make sure you avoid PCR-induced errors by using a PCR-free plasmid prep kit.
- 9. Decrease vector assembly time**
Vector assembly time is a key factor in the efficiency of your assembly. Make sure you decrease vector assembly time by using a vector assembly tool like G-Gate or G-Gate Online.
- 10. Consider droplet-based**
Droplet-based金屬化 is a great way to increase the efficiency of your assembly. Make sure you consider droplet-based金屬化 when planning your reaction.
- 11. Check for common errors**
Common errors can lead to inaccurate results. Make sure you check for common errors like enzyme compatibility, vector compatibility, and buffer compatibility.

Golden Gate Assembly

20+ FRAGMENT ASSEMBLY NOW ACHIEVABLE WITH HIGH EFFICIENCY AND ACCURACY



~~SECRET//NOFORN~~


~~SECRET//NOFORN~~

WIV Bat Coronavirus Collection Efforts

- WIV possesses a large bank of Bat Coronaviruses isolated from various bat species in Yunnan Province China
- Yang et al., 2016
- Hu et al., 2017
- Five-year longitudinal study to isolate Bat Coronaviruses (April 2011 – October 2015)
- Only a few sequences have been published

Table 1. Summary of SARS-CoV detection in bats from a single habitat in Kunming, Yunnan.

Sampling time	Sample type	Sample Numbers			SARS-CoV + bat species (No.)
		Total	CoV +	SARS-CoV +	
April, 2011	anal swab	14	1	1	<i>R. sinicus</i> (1)
October, 2011	anal swab	8	3	3	<i>R. sinicus</i> (3)
May, 2012	anal swab & feces	54	9	4	<i>R. sinicus</i> (4)
September, 2012	feces	39	20	10	<i>R. sinicus</i> (16)
April, 2013	feces	52	21	18	<i>R. ferrugineus</i> (3)
July, 2013	anal swab & feces	115	9	8	<i>R. sinicus</i> (8)
May, 2014	feces	131	8	4	<i>A. stenodermus</i> (3)
October, 2014	anal swab	19	4	4	<i>R. sinicus</i> (1)
May, 2015	feces	145	3	0	<i>R. sinicus</i> (4)
October, 2015	anal swab	25	6	5	<i>R. sinicus</i> (5)
<i>Total</i>		602	84	64	<i>R. (61) A. (3)</i>



- (1) Yunnan
- (2) Guizhou
- (3) Guangxi
- (4) Hong Kong
- (5) Hubei
- (6) Shaanxi
- (7) Hebei
- (8) Jilin


The next page is DIF citing (b)(1) and (b)(3) and is not provided.

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

Isolation Locations for RaTG13 and MP789

- RaTG13/4991 isolated from a cave in Yunnan ProvinceRaTG13/4991 is a RdRp lineage 1 BetaCoV/MP789 was isolated from diseased Pangolins in Guangdong ProvinceMP789 is a RdRp lineage 2 BetaCoV~800 miles separate these two locationsWIV also collected CoV's from Guandong and may have a MP789-related virus in their bank“All the genomic constituents of SARS-CoV including the hypervariable regions S and ORF8 were discovered from different bat SARSr-CoVs in the same cave in Yunnan, with evidence of recombination events detected between these bat SARSr-CoVs...” (Yu et al., 2019)Question: How would a Pangolin RBD from 800 miles away in Guangdong Province recombine into a BatCoV in Yunnan Province?



~~SECRET//NOFORN~~

Postulated WIV Bat-CoV-X Full-length Clone Construction Process

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

Quote from Zeng *et al.*, 2016

- From Materials and Methods, Virus and cells section
“All experiments using live virus was conducted under biosafety level 2 (BSL2) conditions.”

From the first paragraph of the Discussion

“In this study, we have developed a fast and cost-effective method for reverse genetics of coronaviruses by combining two approaches developed by others (29, 30). Our method allows the genomes of coronaviruses to be split into multiple fragments and inserted into a BAC plasmid with a single step. Recombinant viruses can then be efficiently rescued by direct transfection of the BAC construct. As the genomes can be divided into multiple short fragments, mutations can be introduced into individual fragments easily (31). Using this method, we successfully rescued three recombinant viruses derived from SL-CoV WIV1 (rWIV1, rWIV1-DX, and rWIV1-GFP-DX).”

~~SECRET//NOFORN~~

The next page is DIF citing (b)(1) and (b)(3) and is not provided.

~~SECRET//NOFORN~~

WIV SARS-CoV Reverse Genetics System





FIG 1 Strategy for construction of an infectious WIV1 BAC clone. (A) Genomic structure of WIV1. (B) The mutations are indicated under the stars. C1575A was used to ablate a natural BglII site at nucleotide 1571 (▼), and T27327C was used to disrupt a potential T7 stop site. The others were for introducing BglII sites (▲). (C) The WIV1 genome was split into eight contiguous cDNAs (A to G); A, nt 1 to 4387; B, nt 4388 to 8032; C1, nt 8033 to 10561; C2, nt 10562 to 12079; D, nt 12080 to 17017; E, nt 17018 to 22468; F, nt 22469 to 27352; G, nt 27353 to 30309. Unique BglII sites were introduced into the fragments by synonymous mutations to make these fragments capable of undirectional ligation along with native BglII sites in the genome. The original nucleotides are shown above the flanking sequences of corresponding fragments. A poly(A) sequence was added to the 3' terminus of fragment G. A CMV promoter, HDV ribozyme, and BGH transcriptional terminal signal were inserted into the pBeloBAC11 between BamHI and HindIII sites. SacII and AscI sites were introduced between the CMV promoter and ribozyme. Fragments A to G were inserted into the pBAC-CMV plasmid in a single step.

SARS-COV-2 GENOME RESTRICTION MAP

~~SECRET//NOFORN~~



~~SECRET//NOFORN~~

Type IIIS Restriction Enzymes


- BsmBI (Plus strand)
5' - CGTCTC_NNNNN-3'
3' - GCAGAGNNNN - 5' BsaP' (PLNSNN_NGATTC)GC-
- BsmBI (Minus strand)
5' - NNNNNGAGACG-3'
5' BsaI (Minus strand)
- 5' - GGTCTC_{NN}NN-3'
3' - CCAGAGNNNN - 5' SARS-CoV-2_NNNNN_NGA_NCG_N - 3', not have any SacII or AscI restriction sites

(b)(1); (b)(3)(i) USC 3024(i); Sec. 1.4(c); Sec. 1.4(e)

~~SECRET//NOFORN~~


RE-based Fragment Build Option – *BsmBI/BsaI* (4 nt overhangs)

~~SECRET//NOFORN~~



~~SECRET//NOFORN~~

RE-based Fragment Build Option – *BsaI/Esp3I* (Invisible restriction sites)



~~SECRET//NOFORN~~

Xie et al., 2020 SARS-CoV-2 FLC Assembly




Figure 1. Assembly of a Full-Length SARS-CoV-2 Infection cDNA Clone

(A) Genome structure SARS-CoV-2. The open reading frames (ORFs) from the 5' genome are indicated.

(B) Strategy for *in vitro* assembly of an infectious cDNA clone of SARS-CoV-2. The nucleotide sequences and genome locations of the collective overlaps are indicated. The WT full-length (F1) cDNA of SARS-CoV-2 (IC WT) was discontiguously assembled using *in vitro* ligation.

(C) Diagram of the terminal sequences of each cDNA fragment recognized by BamI and EcoRI.

(D) Gel analysis of the seven purified cDNA fragments. Individual fragments (F1-F7) were digested from corresponding plasmid clones and gel-purified. Seven

purified cDNA segments (50–100 ng) were analyzed on a 0.6% native agarose gel. The 1-kb DNA ladder are indicated.

(E) Gel analysis of cDNA ligation products. About 200 ng of purified ligation product was analyzed on a 0.6% native agarose gel. Triangle indicates the FL cDNA product. Circles indicate the intermediate cDNA products.

(F) Gel analysis of RNA transcripts. About 1 μg of *in vitro*-transcribed (IVT) RNAs were analyzed on a 0.6% native agarose gel. DNA ladders are indicated. Because

this is a native agarose gel, the DNA size is not directly correlated to the RNA size. Triangle indicates the genome-long 5' RNA transcript. Circles show the shorter RNA transcripts.

~~SECRET//NOFORN~~


Spike Gene

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

SARS CoV-2 SPIKE GENE

~~SECRET//NOFORN~~



Highest homology to RatG13Pangolin CoV

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

Spike Gene Swapping Using the WIV SARS-CoV Reverse Genetics System

• Hu et al., 2017

swapped out the
WIV1 spike gene for
the spike gene of the
following:Rs4231Rs73
27Rf4075Rs4081Rs40
85Rs4235As6526Rp3






FIG 1 Strategy for construction of an infectious WIV1 BAC clone. (A) Genomic structure of WIV1. (B) The mutations are indicated under the stars. C1575A was used to ablate a natural BpuII site at nucleotide 1571 (▽), and T2752C was used to disrupt a potential T7 stop site. The others were for introducing BpuII sites (▼). (C) The WIV1 genome was split into eight contiguous cDNAs (A to G); A, nt 1 to 4387; B, nt 4388 to 8032; C1, nt 8033 to 10561; C2, nt 10562 to 12079; D, nt 12080 to 17017; E, nt 17018 to 22468; F, nt 22469 to 27352; G, nt 27353 to 30309. Unique BpuII sites were introduced into the fragments by synonymous mutations to make these fragments capable of undirectional ligation along with native BpuII sites in the genome. The original nucleotides are shown above the flanking transcriptional terminal signals. A poly(A) sequence was added to the 3' terminus of fragment G. A CMV promoter, HDV ribozyme, and BGH promoter and ribozyme. Fragments A to G were inserted into the pBAC-CMV plasmid in a single step.

Zeng et al., 2016

Hu et al., 2017

~~SECRET//NOFORN~~


SARS-COV-2 SPIKE GENE SEGMENTS QUERY: PANGOLIN-COV



~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

SARS-CoV-2 SimPLOT



- "A recent study found that a human ACE2-binding ridge in SARS-CoV-2 RBD takes a more compact conformation compared with the SARS-CoV RBD; moreover, several residue changes in SARS-CoV-2 RBD may also enhance its human ACE2-binding affinity [13]. The core residues in RBM which may relate to higher human ACE2-binding affinity than SARS-CoV are 100% identical between SARS-CoV-2 and CoV-Pangolin-2020. Therefore, pangolin-CoV-2020 (CoV-Pangolin/GD) potentially recognizes human ACE2 better than the SARS-CoV."

Difference in Receptor Usage between Severe Acute Respiratory Syndrome (SARS) Coronavirus and SARS-Like Coronavirus of Bat Origin[▼]

Wuze Ren,^{1†} Xuxia Qu,^{2†} Wendong Li,^{1‡} Zhenggang Han,¹ Meng Yu,³ Peng Zhou,¹ Shu-Yi Zhang,⁴ Lin-Fa Wang,^{3*} Hongkui Deng,² and Zhengli Shi^{1*}

State Key Laboratory of Virology, Wuhan Institute of Virology, Chinese Academy of Sciences, Wuhan, China;¹ Key Laboratory of Cell Proliferation and Differentiation of the Ministry of Education, College of Life Sciences, Peking University, Beijing, China;² CSIRO Livestock Industries, Australian Animal Health Laboratory and Australian Biosecurity Cooperative Research Center for Emerging Infectious Diseases, Geelong, Australia;³ and School of Life Science, East China Normal University, Shanghai, China⁴

Received 20 May 2007/Accepted 15 November 2007


Severe acute respiratory syndrome (SARS) is caused by the SARS-associated coronavirus (SARS-CoV), which uses angiotensin-converting enzyme 2 (ACE2) as its receptor for cell entry. A group of SARS-like CoVs (SL-CoVs) has been identified in horseshoe bats. SL-CoVs and SARS-CoVs share identical genome organizations and high sequence identities, with the main exception of the N terminus of the spike protein (S), known to be responsible for receptor binding in CoVs. In this study, we investigated the receptor usage of the SL-CoV S by combining a human immunodeficiency virus-based pseudovirus system with cell lines expressing the ACE2 molecules of human, civet, or horseshoe bat. In addition to full-length S of SL-CoV and SARS-CoV, a series of S chimeras was constructed by inserting different sequences of the SARS-CoV S into the SL-CoV S backbone. Several important observations were made from this study. First, the SL-CoV S was unable to use any of the three ACE2 molecules as its receptor. Second, the SARS-CoV S failed to enter cells expressing the bat ACE2. Third, the chimeric S covering the previously defined receptor-binding domain gained its ability to enter cells via human ACE2, albeit with different efficiencies for different constructs. Fourth, a minimal insert region (amino acids 310 to 518) was found to be sufficient to convert the SL-CoV S from non-ACE2 binding to human ACE2 binding, indicating that the SL-CoV S is largely compatible with SARS-CoV S protein both in structure and in function. The significance of these findings in relation to virus origin, virus recombination, and host switching is discussed.

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

Minimal Receptor Binding Domain Cassette

- WIV scientists previously defined the minimal Receptor Binding Domain cassette that could functionally transfer ACE2 binding capability from one Spike protein to anotherSARS Nucleotide: 930-1554SARS Amino Acid: 310-518Receptor Binding MotifsSARS Nucleotide: 1251-1482SARS Amino Acid: 417-494Homology cut points of SARS-CoV-2 coincide with WIV-identified borders of RBD and RBM



— SARS-CoV-2_WIV02
— Bat CoV RaTG13
— Bat SARSr-CoV ZC45
— Bat SARSr-CoV ZXC21

~~SECRET//NOFORN~~

Furin Cleavage Site

SARS-CoV	CATGTCGACACTTCTATGAGTGGCACATTGGAGCTGGCATTTGTGCTAGTTAC	
1980	H V D T S Y E C D I P I G A G I C A S Y	SARS-CoV-2
CATGTCAACAACTCATATGAGTGTGACATA	CCATTGGTGCGAGGTATATGC	GCTAGTTAT
H V N N S Y E C D I P I G A G I C A S Y	BCoV	RaTG13
CATGTCAATAACTCGTATGAGTGTGACATA	CCTATTGGTGCAGGAATATGCC	CGCTAGTTAT
H V N N S Y E C D I P I G A G I C A S Y	SARS-CoV	CATA
ACGTAGTACTAGCCAAAATCTATGTGGCT	AGTTCTTATT-----	AGTT
2028	H T V S L L R S T S Q K S I	
V ASARS-CoV-2	CAGACTCAGACTAATTCTCCTCGGGCACG	TCAATCCATCATTGCC
2082	Q T Q T N S P R R A R S V A S Q S I I	ABCov RaTG13
CAGACTCAA	ACTAATTCAACGTA	TCTATTATTGCC
T N S	R S V A S Q S I I	A Furin Cleavage Site
		NmeAll Restriction Site

A unique restriction site facilitates identifying the correct *E. coli* clone

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

SARS-CoV-2 Furin Cleavage Site GC Content

- The percent GC of the furin cleavage site insert is 77% compared to ~40% of the surrounding DNA. Contains an NmeAII restriction site. The other CoV's with FCS have a %GC of <55%

Virus	Nucleotide and Amino Acid Sequences																												
	G	I	C	A	S	V	Q	T	Q	T	N	S	P	R	A	R	S	V	A	S	Q	I	I	A	Y	T	M		
SARS-CoV2 (47%)	G	I	C	A	S	V	Q	T	Q	T	N	S	P	R	A	R	S	V	A	S	Q	I	I	A	Y	T	M		
	%GC: 14/35 = 40%																				%GC: 10/13 = 77%								
	C	T	T	G	T	C	T	C	T	G	A	C	A	C	C	T	G	G	T	T	G	T	C	C	A	T	G		
MERS-CoV (54%)	L	C	A	L	P	D	T	P	S	T	L	T	P	R	S	V	R	S	V	P	G	E	M	R	L	A	S	I	A
	%GC: 18/35 = 51%																				%GC: 8/13 = 62%								
	G	G	T	C	A	A	T	C	A	T	T	G	C	T	T	C	G	G	T	T	G	C	A	T	T	G	C		
BatCoV-HKU5 (47%)	G	Q	S	L	C	A	I	P	P	T	T	S	S	R	V	R	R	A	T	S	G	A	S	D	V	F	Q	I	A
	%GC: 15/35 = 43%																				%GC: 7/13 = 54%								
	I	B	V	-B	e	s	d	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t	t		
	Influenza A virus causes a conservative insertion of a poly-A tail at the furin cleavage site by virtue of RNA Polymerase stuttering which adds basic cleavage site by virtue of RNA Polymerase stuttering which adds																				%GC: 13/39 = 33%								


Influenza A virus causes a conservative insertion of a poly-A tail at the furin cleavage site by virtue of RNA Polymerase stuttering which adds basic cleavage site by virtue of RNA Polymerase stuttering which adds preferentially A's and T's – this is not the case with SARS-CoV-2

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

Spike Protein Regions

% Identity:>95



~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

SARS-CoV-2 Spike RBD Alignment: Possible RBM Cassette Insertion

~~SECRET//NOFORN~~

ACE2 Critical Contact ACE2

~~SECRET//NOFORN~~

SARS-CoV-2 RBD vs PCoV MP789 RBD

>REBM


D Y N Y K L P D D F T G C V I A W N S N	MP789	D
GACTATATTATAAACCTGATGATTACAGGTGGTAAAGCTGGATTCTAAC	1305SARSCOV2	
GATTATATTATAATTACCGATGTTAGGCTGGTTATAGCTGGATTCTAAC	1317	
Y N Y K L P D D F T G C V I A W N S N	N L D	
S K V G N Y N Y L Y R L F R K S	MP789	
MP789		
S K V G N Y N Y L Y R L F R K S	1365SARSCOV2	
AACCTGATTCAAGGTTGGTAATTATACTACCTTTATAGTTAGATGTTAGGAAGTC	1377	
L D S K V G N Y N Y L Y R L F R K S	N L K	
P F E R D I S T E I Y Q A G S T P	MP789	
AACCTCAAACCTTTGACGAGACATTCTACAGAAATATACCAAGCTGGTAGCACCC	1425SARSCOV2	
AATCTCAACCTTTGAGAGATATTCACTGAAATCTATCAGGCCGGTAGCACCT	1437	
L K P F E R D I S T E I Y Q A G S T P	C N G	
V E G F N C Y F P L Q S Y G F Q P	MP789	
TGCATGGGTTGAAGGTTTACTGTACTTCCTACATGGTTCCACCC	1485SARSCOV2	
TGTATGGTGTGAAGGTTTATGTACTTCCTACATGGTTCCACCC	1497	
N G V E G F N C Y F P L Q S Y G F Q P	C	
RBM< T N G V G Y Q P Y R V V V L S F E L L H		
MP789		
ACTAATGGTGTGGTACCAACCTATAGAGTAGTAGTATGGTCACTTTAAA		
1545SARSCOV2		
ACTAATGGTGTGGTACCAACCATACAGAGTAGTAGTACTTCTTGGACTCTACAT	1557	
N G V G Y Q P Y R V V V L S F E L L K	T	
RBD< A P A T V C G P K K S T N	MP789	
GCACCTGCTACTGTTGGGACTTAACAGTCCACTAACCTAGTAAACAAATGTGC	1605SARSCOV2	
GCACCAAGCAACTGTTGGACCTAAAGTCTACTATTGGTAAACAAATGTGC	1617	
P A T V C G P K K S T N	A	

- 38 codon differences
- First: 4Second: 0Third: 31First and third: 31 results in an amino acid change
- Pangolin RBD cassette appears to be a codon optimized insert

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

SARS-CoV-2 Spike Appears to be a Chimera




Break points align with those identified by WIV Scientists in
2008 (Ren *et al.*, 2008)

~~SECRET//NOFORN~~

RmYN02 - A Red Herring?

- Zhou et al., 2020 publish paper describing Bat Cov rmyN02Next generation sequencing was done on pooled bat samples to develop two genome sequences – RmyN01 and RmyN02Claim that RmyN02 contains inserted nucleotides at the S1/S2 cleavage siteAssert that the SARS-CoV-2 FCS is therefore of natural originNo virus is available for peer confirmation



~~SECRET//NOFORN~~

Zhou et al., 2020

~~SECRET//NOFORN~~

RmYN02 - A Red Herring?

Virus	Nucleotide Sequences									
SARS-CoV2	G	G	T	A	T	A	T	G	C	T
RaTG13	G	G	A	A	T	A	T	G	C	C
P-CoV MP789	G	G	A	A	T	A	T	G	T	G
RmYN02	G	G	T	G	I	G	I	G	C	A
ZC45	G	G	T	A	T	T	T	G	C	T
ZXC21	G	G	T	A	T	T	T	G	C	T
SARS-CoV	G	G	C	A	T	T	T	G	C	T

~~SECRET//NOFORN~~

HIV Epitopes


Perez, 2020

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

HIV Sequences in the SARS-CoV-2 Spike Gene



Adapted from Perez, 2020

~~SECRET//NOFORN~~

Perez, 2020 Scientific Challenges

- None of the six proposed regions are identical at either the nucleotide or amino acid level with the corresponding HIV/SIV segmentsNone of the six peptides are related to identified immunosuppressive regions of HIV and SIV (Retroviral ISU Domains)The HIV gp41 Immunosuppressive (ISU) Domains sequence is KQLQARILAVERYLKDQQQLLGG - this sequence does not match any of the sixFour of the six regions either perfectly or almost perfectly match corresponding peptides in multiple Pangolin CoVs - Perez did not account for Pangolin genomes in the paperSeveral are only found in Pangolin CoV Spike sequences and not in Bat CoV Spike sequences, indicating that the SARS-CoV-2 Spike NTD region originated from a Pangolin CoV template

The next 2 pages are DIF citing (b)(1) and (b)(3)
and are not provided.

~~SECRET//NOFORN~~

Alternative Scenario

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

Hypothetical Laboratory Origin of SARS-CoV-2

- WIV conducted a longitudinal studies to isolate a large number of bat Coronaviruses from multiple locations in China (2011-2015)WIV Developed Reverse Genetic System, assembled WIV1 full-length infectious clone, and created chimeric viruses exchanging the WIV1 spike gene with the spike gene from other bat Coronaviruses (2015-2017)WIV and other Chinese scientists conduct gain of function studies on SARS, MERS, IBV, and PEDV to insert furin cleavage sites demonstrating increased virulence of the chimeric virusesWIV conducted in vivo and in vitro studies to characterize the bank of bat CoronavirusesWIV conducted the live bat Coronavirus studies under BSL2 conditionsChinese BSL2 and US BSL2 conditions are differentChinese labs have had a history of virus escapes from BSL2 laboratoriesHypothesis: Between 2017 and 2019, WIV created a full-length infectious clone in pBAC-CMV using an unpublished bat Coronavirus genome as template (BatCoVX)Hypothesis: Between 2017 and 2019, WIV created chimeric Bat-CoV-X viruses using the pBAC-CMV-BCoVX backbone and swapping out key cassettes with other bat Coronaviruses (RBD, RBM, etc.) and adding additional features such as a furin cleavage siteHypothesis: In 2018-2019, WIV conducted in vitro and in vivo studies to characterize the BatCoVX chimeric viruses under BSL2 conditionsHypothesis: In mid-2019, one of the not fully characterized Bat-CoV-X chimeric viruses escaped from the WIV facilities and begins infecting civilians in the city of WuhanHypothesis: Starting in mid-2019 through present, WIV and other Chinese laboratories conduct studies to characterize the Chimeric BCoVX virus that escaped (now called SARS-CoV-2)WIV (Zhou et al., 2020) publishes the 2019-nCoV genome sequence showing relatedness to RaTG13 (a previously unpublished genome)BatCoVX likely highly related to RaTG13Hypothesis: Beginning in early 2020, WIV and other government controlled agencies begin to publish obfuscation information to drive the narrative that SARS-CoV-2 is of natural origin and resulted from natural recombinationRaTG13RMYN02pangolin CoVs

~~SECRET//NOFORN~~

CONCLUSION

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~

The next page is DIF citing (b)(1) and (b)(3) and is not provided.

~~SECRET//NOFORN~~

Concluding Points

WIV possesses a bank of Bat Coronavirus isolates WIV has scientists experienced in Coronavirology and Coronavirus Infectious Clone generation WIV Scientists generated chimeric SARS-CoV and Bat CoV Spike genes to identify minimal Spike Receptor Binding Domain cassette that could transfer receptor binding specificity (Ren et al., 2008) WIV possesses an existing and published Coronavirus Reverse Genetics System (Zeng et al., 2016) utilizing their pBAC-CMV plasmid WIV has utilized the pBAC-CMV-WIV1 Full-length clone to generate chimeras with Bat CoV spike genes (Hu et al., 2017) WIV has BSL2/BSL3/BSL4 animal facilities WIV has multiple in vitro assays (apoptosis, IFNB induction, etc.) to characterize their Bat Coronaviruses and chimeric Bat Coronaviruses WIV and other Chinese researchers have conducted Gain of Function studies in SARS, MERS, IBV, and PEDV to add Furin Cleavage Sites to CoV Spike protein The absence of a published progenitor virus for SARS-CoV-2 only indicates that it has not been published, not that it does not exist The genomic sequence of SARS-CoV-2 has Type IIS restriction sites that are consistent with being generated by the Golden Gate Cloning system utilizing the published pBAC-CMV plasmid The SARS-CoV-2 genome has several break points where homology jumps from Bat Coronaviruses to Pangolin Coronaviruses which is consistent with a synthesized chimeric virus The SARS-CoV-2 Spike protein similarity with RaTG13 and Pangolin CoV Spike proteins may also be explained by use of cassettes swapped into the base virus – these break points align with those identified by WIV scientists (Ren et al., 2008) The Pangolin RBD cassette is 100% identical at the amino acid level while the DNA sequence appears to be codon optimized There are no other published SARS lineage Betacoronaviruses that possess a Furin Cleavage Site in their Spike protein (RmYN02 does not have an insertion) and the SARS-CoV-2 FCS does not appear to be inserted via the same mechanism that drives Influenza virus insertions of polybasic cleavage sites Zeng et al., 2016 stated that "All experiments using live virus was conducted under biosafety level 2 (BSL2) conditions" which would make an accidental release of a pathogenic Bat CoV capable of binding human ACE2 more likely A chimeric virus comprised of segments from natural Bat CoV genomes would appear like a recombinant virus

The molecular biology capabilities of WIV and the genome assessment are consistent with the hypothesis that SARS-CoV-2 was a lab-engineered virus that was part of a bank of chimeric viruses in Zhen-Li Shi's laboratory at WIV that escaped from containment

~~SECRET//NOFORN~~


BACK-UP SLIDES

~~SECRET//NOFORN~~

~~SECRET//NOFORN~~


Nucleotide 914 Region

~~SECRET//NOFORN~~




Nucleotide 1312 Region

~~SECRET//NOFORN~~



Nucleotide 1535 Region

~~SECRET//NOFORN~~



~~SECRET//NOFORN~~

SARS-CoV 1980	CATGTCGACACTTCTATGAGTGCAGCATTCCCTATTGGAGCTGGCATTGTGCTAGTTAC
H V N N S Y E C D I P I G A G I C A S Y	SARS-CoV-2
CATGTCAACTCATGAGTGTGACATAACCATTGGTGCA	<u>GAGGTATGCC</u> TAGTTAT
H V N N S Y E C D I P I G A G I C A S Y	RatG13
CATGTCAACTCGTATGAGTGTGACATAACCTATTGGTGCA	GAGGAATATGCCAGTTAT
H V N N S Y E C D I P I G A G I C A S Y	2022
ACGTAGTACTAGCCAAAATCTATTGTGGCT	SARS-CoV
V ASARS-CoV-2	CATA CAGTTCTTATT-----
2082	2028
Q T Q T N S P R R A R S V A S Q S I I	H T V S L L R S T S Q K S I
CAGACTCAA	ABCov RatG13
TA	ACGTAGTGTGGCCAGTCAATCTTATTGCC
Restriction Site	2070
Q T Q	Q T Q
Restriction Site	2070
N	Q T Q
R S V A S Q S I I	Q T Q
Restriction Site	Q T Q
Afuri Cleavage Site	
BspMI Restriction Site	
NmeAI Restriction Site	
BsrDI Restriction Site	

~~SECRET//NOFORN~~