

UNCLASSIFIED

AFMIC Home

Defense Intelligence Agency
DIA SEALAFMIC SEAL

Defense Intelligence Assessment

(U) Infectious Disease Risk Assessment: Iraq

October 2004

DI-1812-IRQ-04

Information Cutoff Date: 07 September 2004

Key Judgments

- AFMIC assesses Iraq as **INTERMEDIATE RISK** for infectious diseases, with an overall disease risk that will adversely impact mission effectiveness unless force health protection measures are implemented.

- **High Risk Diseases:** The main force health protection emphasis should be on these diseases, which are the most likely to degrade operations by affecting a large percentage of personnel, or by causing severe illness in a smaller percentage. High risk diseases are grouped into transmission categories that are prioritized in descending order of risk.

Foodborne And Waterborne Diseases

Diarrhea - bacterial, Hepatitis A,
Typhoid / paratyphoid fever

Vector-borne Diseases

As a group may constitute a high
risk; see table

- **Intermediate Risk Diseases:** These diseases also warrant force protection emphasis. They are less likely to degrade operations because they generally affect smaller numbers of personnel, or cause mild symptoms. Other intermediate risk diseases are those assessed to be present at unknown levels that, under conditions favorable for transmission, could degrade operations. See Detailed Information on Intermediate Risk Diseases.
- **Low Risk Diseases:** Other militarily significant diseases that are likely to have a minimal impact on operational readiness. See List of Diseases of Low Risk or the Summary Table.

Contents

- Key Judgments
- High Risk Diseases
- Intermediate Risk Diseases
- Summary Table
- Appendix

(b)(3):50 USC 3024(i)

(b)(3):10 USC 424;(b)(3):50 USC 3024(i)

(b)(3):50 USC 3024(i)

Foodborne And Waterborne Diseases of High Risk

Overview:

Though improvements are being made, sanitation remains generally poor throughout the country, including major urban areas. Local food and water sources (including ice) are heavily contaminated with pathogenic bacteria, parasites, and viruses to which most US service members have little or no natural immunity.

Disease surveillance is improving since Operation Iraqi Freedom but remains rudimentary. In the years prior to the operation, only a small fraction of disease was identified or reported.

Diarrheal diseases can be expected to temporarily incapacitate a very high percentage of personnel within days if local food, water, or ice is consumed. Hepatitis A and typhoid fever can cause prolonged illness in a smaller percentage.

In addition, though not specifically assessed in this document, viral gastroenteritis (e.g., norovirus) and food poisoning (e.g., *Bacillus cereus*, *Clostridium perfringens*, *Staphylococcus*) may cause significant outbreaks.

The diseases of high risk are listed first, in descending order of expected impact. Diseases of intermediate risk, with a lower or unknown likelihood to degrade operations, are listed alphabetically in tabular form.

Diarrhea - bacterial

(b)(3):10 USC 424;(b)(3):50 USC 3024(i)

Risk Assessment:

- An operationally significant attack rate [] could occur among personnel consuming local food, water, or ice. Field conditions (including lack of handwashing and primitive sanitation) may facilitate person-to-person spread and epidemics.
- Typically mild disease treated in outpatient setting; recovery and return to duty in less than 72 hours with appropriate therapy.

Risk Period: Year-round

Risk Distribution: Countrywide (including urban areas)

Typical Incubation Period: 1 to 3 days

Epidemiology Comments: In general, bacterial agents such as enterotoxigenic *E. coli*, *Campylobacter*, *Shigella*, and *Salmonella* are the most common causes of "traveler's diarrhea" wherever sanitary conditions are significantly below US standards.

(b)(3):10 USC 424;(b)(3):50 USC 3024(i)

Hepatitis A

Risk Assessment:

Page 5 is redacted in full IAW b3 exemption. Page not included.

- Risk Period:** Year-round

Typical Incubation Period: 28 to 30 days (maximum range: 15 to 50 days)

(b)(3):10 USC 424;(b)(3):50 USC 3024(i)

(b)(3):10 USC 424;(b)(3):50 USC 3024(i)

- A small number of cases [] could occur among unvaccinated personnel consuming local food, water, or ice.
- Debilitating febrile illness typically requiring 1-7 days of inpatient care, followed by return to duty.

Risk Distribution: Countrywide (including urban areas); risk is elevated in populated areas with poor sanitation.

Epidemiology Comments: Typhoid and paratyphoid are clinically similar, and in endemic areas typhoid typically accounts for 90 percent of cases. Asymptomatic carriers are common with typhoid and contribute to sustained transmission. In countries where hygiene and sanitation are poor or nonexistent, adult cases and outbreaks are rare because of immunity acquired in infancy or early childhood. In countries with a mixture of primitive and modern sanitation and hygiene, outbreaks of typhoid fever occur and may involve all age groups.

(b)(3):10 USC 424;(b)(3):50 USC 3024(i)

Page 05 of 14

Withheld pursuant to exemption

(b)(3):10 USC 424;(b)(3):50 USC 3024(i)

of the Freedom of Information and Privacy Act

Foodborne And Waterborne Diseases of Intermediate Risk

Click on disease for additional information	(b)(3):10 USC 424;(b)(3):50 USC 3024 (i)
Brucellosis	
Diarrhea - cholera	
Diarrhea - protozoal	
Hepatitis E	

Vector-borne Diseases of Intermediate Risk

The seasonality and geographic distribution of vector-borne diseases may vary considerably. Click on the disease for details

Overview:

During warmer months (typically April through November), ecological conditions support populations of arthropod vectors, including mosquitoes, ticks, and sand flies, with variable rates of disease transmission.

Plasmodium vivax malaria occurs at low levels in some rural areas. In addition, a variety of other vector-borne diseases occur at low or unknown levels, which as a group may constitute a significant risk. Personnel exposed to mosquitoes, ticks, sand flies, or other biting vectors are at risk during day or night.

Click on disease for additional information	(b)(3):10 USC 424;(b)(3):50 USC 3024 (i)
Crimean-Congo hemorrhagic fever	
Leishmaniasis - cutaneous	
Leishmaniasis - visceral	
Malaria	
Plague	
Rickettsioses, tickborne (spotted fever group)	
Sand fly fever	
West Nile fever	

Other Diseases of Intermediate Risk

Click on disease for additional information	(b)(3):10 USC 424;(b)(3):50 USC 3024(i)
Sexually Transmitted Diseases	
Gonorrhea / chlamydia	
HIV/AIDS	
Hepatitis B	
Water-contact Diseases	
Leptospirosis	
Schistosomiasis	
Respiratory Diseases	
Tuberculosis	
Tuberculin skin test (TST) conversion rates may be elevated over baseline for personnel with prolonged close exposure to local populations	
Animal-contact Diseases	
Anthrax	
Q fever	
Rabies	

List of Diseases of Low Risk

Meningococcal meningitis
 Sindbis (Ockelbo) virus
 Typhus - murine (fleaborne)

Administrative Notes

(b)(3):10 USC 424

This report contains information as of 07 September 2004. This document, published under the auspices of the Department of Defense Intelligence Production Program (DoDIPP), reflects the Defense Intelligence Production Community position. The Defense Intelligence Agency's Armed Forces Medical Intelligence Center produced it as the designated DoDIPP producer for this subject.

This publication supersedes Infectious Disease Risk Assessment: Iraq, DI-1812-IRQ-04, dated January 2004, which should be destroyed.

(b)(3):10 USC 424;(b)(3):50 USC 3024(i)

UNCLASSIFIED

AFMIC Home Page

Limited Value ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 High Value

Name:

Email:

Organization:

Phone:

Comments:

UNCLASSIFIED

Defense Intelligence Agency

Defense Intelligence Assessment

(U) Infectious Disease Risk Assessment: Iraq

October 2004

(b)(2)

Information Cutoff Date: 07 September 2004

Key Judgments

- AFMIC assesses Iraq as **INTERMEDIATE RISK** for infectious diseases, with an overall disease risk that will adversely impact mission effectiveness unless force health protection measures are implemented.

(b)(1), Sec. 1.4(e)

- **High Risk Diseases:** The main force health protection emphasis should be on these diseases, which are the most likely to degrade operations by affecting a large percentage of personnel, or by causing severe illness in a smaller percentage. High risk diseases are grouped into transmission categories that are prioritized in descending order of risk.

Foodborne And
Waterborne Diseases

Diarrhea - bacterial, Hepatitis A, Typhoid
/ paratyphoid fever

Vector-borne Diseases

As a group may constitute a high risk; see
table

- **Intermediate Risk Diseases:** These diseases also warrant force protection emphasis. They are less likely to degrade operations because they generally affect smaller numbers of personnel, or cause mild symptoms. Other intermediate risk diseases are those assessed to be present at unknown levels that, under conditions favorable for transmission, could degrade operations.

(b)(1), Sec. 1.4(c)

(b)(1), Sec. 1.4(c)

- **Low Risk Diseases:** Other militarily significant diseases that are likely to have a minimal impact on operational readiness. See List of Diseases of Low Risk or the Summary Table.

(b)(1), Sec. 1.4(c)

Contents

- Key Judgments
- High Risk Diseases
- Intermediate Risk Diseases
- Summary Table
- Appendix

(b)(1), Sec. 1.4(e)

Foodborne And Waterborne Diseases of High Risk

Overview:

Though improvements are being made, sanitation remains generally poor throughout the country, including major urban areas. Local food and water sources (including ice) are heavily contaminated with pathogenic bacteria, parasites, and viruses to which most US service members have little or no natural immunity.

Disease surveillance is improving since Operation Iraqi Freedom but remains rudimentary. In the years prior to the operation, only a small fraction of disease was identified or reported.

Diarrheal diseases can be expected to temporarily incapacitate a very high percentage of personnel within days if local food, water, or ice is consumed. Hepatitis A and typhoid fever can cause prolonged illness in a smaller percentage.

In addition, though not specifically assessed in this document, viral gastroenteritis (e.g., norovirus) and food poisoning (e.g., *Bacillus cereus*, *Clostridium perfringens*, *Staphylococcus*) may cause significant outbreaks.

The diseases of high risk are listed first, in descending order of expected impact. Diseases of intermediate risk, with a lower or unknown likelihood to degrade operations, are listed alphabetically in tabular form.

Diarrhea - bacterial

Risk Assessment:

- An operationally significant attack rate (b)(1), Sec. 1.4(c) could occur among personnel consuming local food, water, or ice. Field conditions (including lack of handwashing and primitive sanitation) may facilitate person-to-person spread and epidemics.
- Typically mild disease treated in outpatient setting; recovery and return to duty in less than 72 hours with appropriate therapy.

Risk Period: Year-round

Risk Distribution: Countrywide (including urban areas)

Typical Incubation Period: 1 to 3 days

Epidemiology Comments: In general, bacterial agents such as enterotoxigenic *E. coli*, *Campylobacter*, *Shigella*, and *Salmonella* are the most common causes of "traveler's diarrhea" wherever sanitary conditions are significantly below US standards.

(b)(1), Sec. 1.4(c)

Hepatitis A

Risk Assessment:

- An operationally significant attack rate (b)(1), Sec. 1.4(c) could occur among unvaccinated personnel consuming local food, water, or ice. Field conditions (including lack of

handwashing and primitive sanitation) may facilitate person-to-person spread and epidemics.

- Typical case involves 1 to 3 weeks of debilitating symptoms, sometimes initially requiring inpatient care; recovery and return to duty may require a month or more.

Risk Period: Year-round

Risk Distribution: Countrywide (including urban areas)

Typical Incubation Period: 28 to 30 days (maximum range: 15 to 50 days)

Epidemiology Comments: In non-indigenous personnel, hepatitis A typically occurs after consumption of contaminated food or water. Infection also may occur through direct fecal-oral transmission under conditions of poor hygiene and sanitation. Childhood hepatitis A infections tend to be asymptomatic or have mild symptoms, while adults typically develop jaundice and other symptoms. In areas where high levels of hepatitis A virus circulate, the number of reported cases in the local population is usually low because asymptomatic childhood infections confer lifelong immunity.

(b)(1),Sec. 1.4(c)

Typhoid / paratyphoid fever

Risk Assessment:

- A small number of cases (b)(1),Sec. 1.4(c) could occur among unvaccinated personnel consuming local food, water, or ice.
- Debilitating febrile illness typically requiring 1-7 days of inpatient care, followed by return to duty.

Risk Period: Year-round; risk is elevated during warmer months.

Risk Distribution: Countrywide (including urban areas); risk is elevated in populated areas with poor sanitation.

Typical Incubation Period: 8 to 14 days (maximum range: 3 to 30 days)

Epidemiology Comments: Typhoid and paratyphoid are clinically similar, and in endemic areas typhoid typically accounts for 90 percent of cases. Asymptomatic carriers are common with typhoid and contribute to sustained transmission. In countries where hygiene and sanitation are poor or nonexistent, adult cases and outbreaks are rare because of immunity acquired in infancy or early childhood. In countries with a mixture of primitive and modern sanitation and hygiene, outbreaks of typhoid fever occur and may involve all age groups.

(b)(1),Sec. 1.4(c)

(b)(1),Sec. 1.4(c)

Foodborne And Waterborne Diseases of Intermediate Risk

Click on disease for additional information	(b)(1),Sec. 1.4(c)
---	--------------------

<u>Brucellosis</u>	(b)(1), Sec. 1.4(c)
<u>Diarrhea - cholera</u>	
<u>Diarrhea - protozoal</u>	
<u>Hepatitis E</u>	

Vector-borne Diseases of Intermediate Risk

The seasonality and geographic distribution of vector-borne diseases may vary considerably. Click on the disease for details

Overview:

During warmer months (typically April through November), ecological conditions support populations of arthropod vectors, including mosquitoes, ticks, and sand flies, with variable rates of disease transmission.

Plasmodium vivax malaria occurs at low levels in some rural areas. In addition, a variety of other vector-borne diseases occur at low or unknown levels, which as a group may constitute a significant risk. Personnel exposed to mosquitoes, ticks, sand flies, or other biting vectors are at risk during day or night.

Click on disease for additional information	(b)(1), Sec. 1.4(c)
<u>Crimean-Congo hemorrhagic fever</u>	
<u>Leishmaniasis - cutaneous</u>	
<u>Leishmaniasis - visceral</u>	
<u>Malaria</u>	
<u>Plague</u>	
<u>Rickettsioses, tickborne (spotted fever group)</u>	
<u>Sand fly fever</u>	
<u>West Nile fever</u>	

Other Diseases of Intermediate Risk

Click on disease for additional information	(b)(1), Sec. 1.4(c)
Sexually Transmitted Diseases	
<u>Gonorrhea / chlamydia</u>	
<u>HIV/AIDS</u>	
<u>Hepatitis B</u>	
Water-contact Diseases	
<u>Leptospirosis</u>	
<u>Schistosomiasis</u>	
Respiratory Diseases	

Tuberculosis Tuberculin skin test (TST) conversion rates may be elevated over baseline for personnel with prolonged close exposure to local populations			
Animal-contact Diseases			
<u>Anthrax</u>		(b)(1), Sec. 1.4(c)	
<u>Q fever</u>			
<u>Rabies</u>			

List of Diseases of Low Risk

Meningococcal meningitis
Sindbis (Ockelbo) virus
Typhus - murine (fleaborne)

Administrative Notes

(b)(6)

(b)(2)

UNCLASSIFIED



Limited Value ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 High Value

Name:

Email:

Organization:

Phone:

Comments: